



**BRITISH HYDROLOGICAL SOCIETY**

**EIGHTH  
NATIONAL  
HYDROLOGY  
SYMPOSIUM**

**THE UNIVERSITY OF  
BIRMINGHAM**

**8-11 SEPTEMBER  
2002**

# CONTENTS

|  |  | <i>Page s</i> |
|--|--|---------------|
| <i>Agnew, C</i>  | Drought and desiccation – the need for new approaches  | 1             |
| <i>Archer, D.</i>  | Flood risk evaluation in the Karakoram Himalaya: implications of non-homogeneity and non-stationarity  | 7             |
| <i>Beasley, G. and Kneale, P.E.</i>  | Contamination risk and in-stream ecological stress: metals, PAHs and macroinvertebrates  | 13            |
| <i>Beaudin, I., White, S.M., Hollis, J., Hallett, S., Worrall, F. and Whelan, M.</i> | The use of SWAT-2000 for diffuse source contaminant modelling at the regional scale across Europe  | 21            |
| <i>Beran, M.</i>   | What price the Precautionary Principle: flood estimation and climatic change   | 29            |
| <i>Booker, D.J., Durr, C.S., Acreman, M.C. and Shamseldin, A.</i>                    | Modelling the effects of altered flow regimes on physical habitat in urban rivers  | 39            |
| <i>Bower, D., Hannah, D.M and McGregor, G.R.</i>                                     | A tool for assessing the climatic sensitivity of river flow regimes  | 49            |
| <i>Charlton, R.A., Moore, S., Fealy, R. and Sweeney, J.</i>                          | Climate change and water resources in Ireland: predictions made at a high spatial resolution using downscaled GCM data and hydrological modelling techniques | 61            |
| <i>Collins, D.N., MacDonald, O.G., Reynolds, A.C. and Rees, H.G.</i>                 | Climatic variation, mass balance and runoff from European glacierised mountain basins  | 67            |
| <i>Dawson, C.W.</i>  | An international comparative study of artificial neural network techniques for river stage forecasting   | 75            |
| <i>Fowler, H.J., Kilsby, C.G. and O'Connell, P.E.</i>                                | Modelling the impact of climate change on the reliability of the Yorkshire water supply network  | 83            |
| <i>Gosling, R.D., Black, A.R. and Brock, B.W.</i>                                    | High snow melt rates in the Cairngorms, N.E. Scotland  | 91            |
| <i>Holmes, M.G.R. and Young, A.R.</i>  | Estimating seasonal low flow statistics in ungauged catchments   | 97            |
| <i>Houghton-Carr, H., Fry, M.G.R., McCartney, M. and Folwell, S.</i>                 | Drought and drought management in southern Africa  | 103           |
| <i>Kannan, N, White, S., Worrall, F., Pendlington, D. and Groves, S.</i>             | Modelling of pesticide runoff from the Colworth experimental catchment, Bedfordshire   | 109           |
| <i>Kilsby, C.G. and Fowler, H.J.</i>   | Changing UK extreme rainfall and implications for design   | 113           |

|   |   |     |
|---|---|-----|
| <i>Littlewood, I.G.</i>   | An improved procedure for identifying quick and slow response unit hydrographs for regionalisation of high and low flows                | 121 |
| <i>Macdonald, N., Black, A.R. and Werritty, A.</i>  | Use of historical data in flood frequency estimation for the River Ouse at York, UK   | 127 |
| <i>Malcolm, I.A., Soulsby, C., Youngson, A.F., Hannah, D., McLaren, I.S. and Thorne, A.</i> | Influence of flow extremes on the hyporheic zone of salmon spawning streams   | 135 |
| <i>McIntyre, N.R., Wheater, H.S. and Borrman, A.J.</i>                                      | Multiple objective evaluation of risk in water resource planning  | 143 |
| <i>Midgley, P.</i>  | Emergency flood planning: the hydrologist's pivotal role in construction of the Lavant Rife   | 151 |
| <i>Parsons, S.B.</i>  | Development of a flood management strategy for Tajikistan   | 157 |
| <i>Pearce, J.</i>   | Hydrologist experiences in times of flood: Sussex 2000  | 163 |
| <i>Reed, D.W.</i>   | Why analyse inter-site dependence in river flooding?  | 169 |
| <i>Robbins, G.L. and Collier, C.G.</i>  | On the assimilation of microwave link attenuation measurements of precipitation into an urban drainage network model                    | 175 |
| <i>Rodgers, P., Soulsby, C., Petry, J., Gibbins, C., Malcolm, I. and Dunn, S.M.</i>         | A tracer-based investigation of groundwater - surface water interactions in a braided river: a case study of the river Feshie, Scotland | 183 |
| <i>Tindall, C.I., Moore, R.V. and George, T.B.</i>  | Data management for the Lowland Catchment Research (LOCAR) Programme  | 191 |
| <i>Ward, R.</i>   | The emergence of British hydrology: reflections after 40 years  | 197 |
| <i>Wheater, H.S., Boxall, S. and Wagener, T.</i>  | Regionalisation of rainfall-runoff models: an application to the Thames Basin   | 199 |
| <i>Wood, P.J. and Agnew, M.D.</i>   | The role of extreme flows in shaping macroinvertebrate communities within groundwater dominated rivers                                  | 207 |
| <i>Zaidman, M., Keller, V. and Young, A.R.</i>  | Trends in streamflow drought occurrence in the UK   | 213 |
| <i>Zhang, J. and Hall, M.J.</i>   | Regional analysis of floods: a case study from south-east China   | 221 |